

## Cervical Screening results and recommendations table



HPV Test Result	Cytology Pattern (where applicable)	Code	Management Recommendation	Rationale/ Recommendation
Not Detected/ Negative	N/A	R1	Screening completed	<ul> <li>Woman is aged 61 or over at date of test.</li> <li>Discharged from colposcopy with a recommendation of nofurther screening.</li> <li>First test at completion of increased surveillance post colposcopy and woman is aged 66 or over at date of test.</li> <li>Woman is aged 64 or over at date of test and has been on annual surveillance for reason of renal dialysis / post-transplant or HIV positive.</li> </ul>
		R3	1 year recall	<ul> <li>Renal dialysis</li> <li>Pre / post-transplant</li> <li>Living with HIV</li> <li>If increased surveillance is indicated as per colposcopy discharge (see note 1 below)</li> </ul>
		R2a	3 year recall	<ul> <li>If aged between 25 and 29 years with no requirement for or completed increased surveillance.</li> <li>First test at completion of increased surveillance post colposcopy - women under 66 years at date of test</li> </ul>
		R2b	5 year recall	<ul> <li>If aged between 30 - 60 years at test date with no requirement for increased surveillance (women must attend for one test at 3 year interval before moving to 5 year recall on completion of increased surveillance post colposcopy if under 66 years at date of test).</li> </ul>
Detected/ Positive	P1 (Unsatisfactory)	R6	3-month repeat	Repeat 3 months
		R7	Refer to colposcopy	<ul> <li>3 consecutive unsatisfactory screening test results</li> <li>Any 3 screening test results that are not normal in previous 10 years &amp; woman has not had colposcopy.</li> <li>Renal dialysis, pre / post-transplant, living with HIV</li> </ul>
	P2 (No abnormality	R3	1 year recall	First HPV positive and cytology NAD, repeat screening test in 1 year (includes women on post colposcopy surveillance)
	detected)	R7	Refer to colposcopy	<ul> <li>Second consecutive HPV positive and cytology NAD taken in non-colposcopy setting. See note 2 below.</li> <li>Living with HIV</li> <li>Renal dialysis</li> <li>Pre / post-transplant</li> </ul>
	P3a+ (ASCUS or worse)	R7	Refer to colposcopy	Any HPV positive result with abnormal cytology
Indeterminate HPV result	N/A	R6	3-month repeat	First or second indeterminate screening test result
		R7	Refer to colposcopy	3 consecutive indeterminate screening test results Any 3 screening test results that are not normal in previous 10 years & woman has not had colposcopy Renal dialysis, pre / post-transplant, living with HIV
Test not Processed	N/A	R6	3-month repeat	Repeat 3 months

NOTES:

1. Women discharged from colposcopy pre-March 2020 for cytology screening who receive a HPV negative result do not require continued annual surveillance unless there is a history of invasive cervical cancer/CGIN/AIS/SMILE. Women discharged after March 2020 should attend for their recommended number of post colposcopy HPV tests.

- 2. Disregard intervening cytology only, indeterminate, unsatisfactory result.
- 3. When current clinical details record Post Coital Bleeding (PCB) / Intermenstrual Bleeding (IMB) / Post-Menopausal Bleeding (PMB) or it is recorded on the screening form that the cervix is suspicious, it is recommended to refer to a gynaecological specialist.
- 4. Where there is a cytology result: If there are endometrial cells present and woman is over 45 years it is recommended to refer to a gynaecological specialist.
- 5. Where there is a cytology result and a previous history of treatment for glandular abnormality: Report as UNSAT if TZ cells are not present.

